




Neural Network for Coated/Uncoated Hydrosol Particles

Grant Sims^{1,*}, Eduard Chemyakin^{1,2}, James Allen³, Jacek Chowdhary^{4,5}, Brian Cairns⁴, Snorre Stamnes^{1,*}



¹NASA Langley Research Center, ²Science Systems and Applications, Inc., ³University of Hawaii, ⁴NASA Goddard Institute for Space Studies (GISS), ⁵Department of Applied Physics and Applied Mathematics, Columbia University

Why Neural Network

- ❖ Accurately reproduce inherent optical properties (IOPs) for coated/uncoated hydrosol particles while reducing required disk space by 100x
- ❖ Provide IOP properties at least 100x faster than the advanced look-up-table

Network Information


- ❖ Coated/uncoated hydrosol particle network:
 - Size: 150 MB
 - Speed: ~ 5,000-10,000 cases/second
- ❖ Input parameters: wavelength, real and imaginary components of the core and shell, (imaginary shell consists of 650 values), effective radius, effective variance
- ❖ Output parameters: extinction coefficient, scattering coefficient, absorption coefficient, lidar backscatter coefficient, bbp, asymmetry parameter, scattering matrix

References

Chemyakin et al., 2021, Improved Lorenz-Mie Look-Up Table for Lidar and Polarimeter Retrievals. Frontiers in Remote Sensing, 2, p.711106.

Network Structure

- ❖ Network structured to take advantage of interdependencies between IOPs (Chemyakin et al., 2021)



- ❖ Uses a custom cubic activation function for output layers in order to produce more accurate results through a wider range of output values

Network Performance

